

PROLIVO™

300SC FUNGICIDE

Put an End to Powdery Mildew

Introducing Prolivo™ 300SC fungicide. A new powdery mildew fungicide for grapes from Summit Agro USA.

Now available for use in wine grapes, table grapes and raisins, Prolivo protects your grapes in multiple ways.

- Excellent preventative activity. Prolivo stops powdery mildew at the onset of the disease epidemic.
- Translaminar activity providing protection to the untreated side of the leaf.
- Through redistribution on the leaf surface and vapor activity, Prolivo offers additional disease protection to untreated leaf and fruit surfaces.
- Prolivo inhibits both lesion formation and prevents sporulation.
- Prolivo has demonstrated excellent rainfastness.

You can count on Prolivo to protect your crop, enhancing yields and quality.


Used as a preventative treatment, apply Prolivo beginning with initial flowering or when conditions are favorable for disease development. Prolivo can be used at rates of 4–5 fluid ounces per acre on a 14 day spray schedule dependent on disease conditions.

Prolivo activity can be improved with the addition of an organosilicant, non-ionic, or mixture adjuvant when used at the rates specified on the adjuvant label.

Make Prolivo a key part of your powdery mildew disease control program today.

Prolivo 300SC Fungicide, a group U8 fungicide, provides preventative disease control of powdery mildew in Caneberry Crop Subgroup 13-07A; Bushberry Crop Subgroup 13-07B; Small Fruit Vine Climbing Crop Subgroup 13-07D; Low Growing Berry Crop Subgroup 13-07G, (except cranberry) and Cucurbit vegetables (Crop Group 9). Active ingredient: Pyriofenone. Distributed through Helena Chemical Company, and Tenkoz member companies.

Always read and follow label directions.

Prolivo is a trademark of Ishihara Sangyo Kaisha, Ltd. 
© 2017 Summit Agro USA LLC. All rights reserved.

Registered crops and disease targets:



Grapes

Powdery Mildew



Cucurbits

Powdery Mildew




Berries

(including strawberries
and other berry crops)
Powdery Mildew

**Ask your SummitAgro
representative for a full
listing of all crop uses and
diseases controlled.**

Grape Disease Control



Evaluation of Fungicides for control of Powdery Mildew in Grapes

	MILDEW ON LEAVES		MILDEW ON BUNCHES	
	% incidence	% severity	% incidence	% severity
Untreated	100a	58a	100a	35a
Prolivo 4 fl oz	1e	0f	0e	0d
Vivando 15 fl oz	0e	0f	0e	0d
Luna Exp. 6 fl oz	1e	0f	0e	0d
Rally 5 oz	78bc	5d	57c	1.5 vc
Topguard 10 fl oz	46cd	1.1e	26d	0.6c
Quintec 4 fl oz	95a	12.4bc	66b	1.7b

2011-BSF303 Dr. Wayne Wilcox, Dept. of Plant Pathology
Cornell University, Geneva, NY. Cultivar Chardonnay

	MILDEW ON LEAVES		MILDEW ON BUNCHES	
	% incidence	% severity	% incidence	% severity
Untreated	92.5	19.5	100	65
Prolivo 4 fl oz	6	.5	59	11
Procure 8 fl oz alt Quintec 4 fl oz	26	1.7	69	15
Topguard	13.5	1.4	62	12
LSD	14.8	2.4	28.7	12.4

2010-BSF302 Nelson & Grove, Washington State Univ.
Prosser, WA, White Riesling variety.

Prolivo is a trademark of Ishihara Sangyo Kaisha, Ltd.; Luna is a trademark of Bayer; Rhyme is a trademark of FMC; Topguard is a trademark of Cheminova; Mettle is a trademark of Isagro USA Inc.; Rally and Quintec are trademarks of Dow Agrosciences LLC; Vivando is a trademark of BASF; Procure is a registered trademark of MacDermid Agricultural Solutions, Inc.

Applying PROLIVO

CROP	RATE	USE INSTRUCTIONS FOR APPLYING PROLIVO
Caneberry Crop Subgroup 13-07A; Bushberry Crop Subgroup 13-07B; Small Fruit Vine Climbing Crop Subgroup 13-07D; Low Growing Berry Crop Subgroup 13-07G, (except cranberry).	4-5 fl. oz.	<ul style="list-style-type: none"> For use on all types of grapes (wine, table, raisin and juice). Applications to grape and similar vine climbing fruits should be made on a 14 day interval. Applications to strawberries and other berry crops should be made on a 7-10 day interval. For powdery mildew control, begin fungicide applications preventatively and continue as needed as per instructions above. Use the low rate and long interval as disease preventative sprays or when disease conditions are low. Increase to highest rate and shortest interval under moderate to heavy disease pressure.
Cucurbit Vegetable Group (9)	4-5 fl. oz.	<ul style="list-style-type: none"> For powdery mildew control, make fungicide applications on a 7-10 day schedule beginning with initial flowering or when disease conditions are favorable for disease development, but prior to disease development. Use the low rate and long interval as disease preventative sprays or when disease conditions are low. Increase to highest rate and shortest interval under moderate to heavy disease pressure.