

# HARVANTA™

## 50SL INSECTICIDE

### New broad-spectrum diamide insecticide offers an unmatched range of performance benefits

Now vegetable growers have an exceptional new option for effective control of their key insect problems: HARVANTA™ 50SL insecticide. Driven by CYCLAPRYN®, a new active ingredient, HARVANTA is a new IRAC Group 28 diamide insecticide that provides a long list of compelling reasons why it should be your first choice for maximizing crop yield and quality;

- Broad-spectrum insect control across many key crops and insects
- Utilizes less active ingredient per acre compared to other Group 28 insecticides
- Excellent partner for Integrated Pest Management (IPM) programs
- Fast acting on contact and ingestion with long residual control. Ingestion or direct contact with the treated leaf leads to anti-feeding, paralysis and the ultimate death of the target insect pest
- Larvicidal and adulticidal activity separating it from other diamide insecticides
- Excellent efficacy against Lepidopteran pests across crop groups when compared to market standards and other Group 28 insecticides
- Effectively controls many insect pests resistant to organophosphate, pyrethroid and carbamate insecticides
- Valuable tool for Insect Resistance Management (IRM) programs

**HARVANTA 50SL insecticide driven by CYCLAPRYN — it's just what you need for better insect management in your vegetable crops.**

### Registered crops and key insect targets:



#### Cucurbits

- Cucumber beetle
- Squash bugs
- Caterpillars/moths
- Cotton/melon aphid



#### Fruiting Vegetables

- Whitefly
- Thrips
- Cutworms/hornworms
- Fruitworms
- Pepper Weevil



#### Brassica (cole) Vegetables

- Diamondback moth
- Imported cabbage worm
- Armyworms
- Flea beetles



#### Leafy (non-brassica) Vegetables

- Diamondback moth
- Armyworms
- Flea beetles

*For a complete list of all controlled insects as well as use instructions, refer to the back of this sheet.*

# Applying HARVANTA™

Performance is enhanced when used with an effective adjuvant. For best results, use the high labeled rate when targeting stink bug and squash bug nymphs. For best results controlling aphids use with an effective adjuvant.

Crop & Rate	Insects	Additional Use Information
<b>Cucurbit Vegetables</b> (Crop group 9)  10.9 to 16.4 fl oz per acre	Beet armyworm Cabbage looper Melon worm Pickleworm Southern armyworm Fall armyworm Striped cucumber beetle Leafminers ( <i>Liriomyza</i> species)	Flea beetles Western yellowstriped armyworm Western flower thrips Onion thrips Whiteflies* Stink bug spp.* Squash bug* Cotton (melon) aphid
<b>Fruiting Vegetables</b> (Crop groups 8-10)  10.9 to 16.4 fl oz per acre	Beet armyworm Cabbage looper Colorado potato beetle Fall armyworm Southern armyworm Tomato fruitworm Tomato hornworm Tomato pinworm Tomato psyllid	Pepper weevil European corn borer Yellow striped armyworm Leafminers ( <i>Liriomyza</i> species) Western flower thrips Whiteflies* Stink bug spp.* Cotton (melon) aphid
<b>Brassica (cole) Leafy Vegetables</b> (Crop groups 5-16)  10.9 to 16.4 fl oz per acre	Beet armyworm Cabbage looper Corn earworm Cross striped cabbage moth Diamondback moth Imported cabbageworm Western yellowstriped armyworm	Flea beetles Leafminers ( <i>Liriomyza</i> species) Western flower thrips Whiteflies* Stink bug spp.* Cotton (melon) aphid
<b>Leafy Vegetables (non-brassica)</b> (Crop groups 4-16)  10.9 to 16.4 fl oz per acre	Beet armyworm Cabbage looper Corn earworm Diamondback moth Flea beetles Fall armyworm	Western yellowstriped armyworm Leafminers ( <i>Liriomyza</i> species) Western flower thrips Whiteflies* Stink bug spp.* Cotton (melon) aphid

Refer to label for complete use information and restrictions. Always follow label directions. \* Suppression only. Use in conjunction with an effective control program.

To see how new HARVANTA, driven by CYCLAPRYN, can help improve your vegetable production, contact your Tenkoz member company or Helena Chemical Company representative, or visit [www.summitagro-usa.com/products](http://www.summitagro-usa.com/products).

**HARVANTA 50SL Insecticide crop import tolerances are in the process of being established. Confirm that tolerances required for the export of your crop have been obtained prior to use.**

**RESISTANCE MANAGEMENT:** Do not apply HARVANTA or other Group 28 insecticide more than three times within a single generation of insect pest(s) on a crop.

## RESTRICTIONS:

- Do not exceed 16.4 fl. oz. per application
- Do not apply more than 65.6 fl. oz./acre/year
- Do not exceed 5 day minimum interval between treatments
- Pre-Harvest Interval (PHI) is 1 day
- Restricted Entry Interval (REI) is 4 hours

To learn more, visit us at [www.summitagro-usa.com](http://www.summitagro-usa.com), or call us at 984-260-0407.

Harvanta insecticide is sold exclusively through Helena Chemical Company and Tenkoz member companies.

HARVANTA and CYCLAPRYN are trademarks or registered trademarks of Ishihara Sangyo Kaisha, LTD © 2017 Summit Agro USA LLC. All rights reserved.