



EPA Reg. No. 71512-31-88783

**Directions for Use on Almonds to Control  
Scab, Alternaria Leaf Spot, and Suppress Hull Rot  
For use in California**

**PRODUCT BULLETIN**

**FIFRA Section 2(ee) Recommendation:** This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**APPLICATION DIRECTIONS**

Crop	Diseases	Use Rate Fl. Oz. Product Per Acre	Instructions
Almond	Scab ( <i>Cladosporium carpophilum</i> )  Alternaria leaf spot ( <i>Alternaria alternata</i> )  Suppression: Hull rot ( <i>Monilinia</i> spp., <i>Rhizopus</i> spp.)	12.5 to 15 fl oz	<p><b>Application Instructions:</b> Initiate applications when conditions are favorable for disease development and continue on a 7-14 day interval. If disease pressure is severe use the higher rate and shortest interval.</p> <p>Apply this product in sufficient water to obtain adequate coverage of the foliage. Spray volume will usually be 50 to 100 gallons per acre for dilute sprays and 5 to 10 gallons per acre for aerial sprays. For aerial applications, apply this product in a minimum of 5 gallons of water per acre. For dilute sprays, if higher spray volumes are desired for improved coverage, do not exceed the maximum rate of 15 fl oz of product per acre.</p> <p><b>Resistance Management:</b> Do not make more than 2 sequential applications of this product or other <b>Group 3</b> or <b>Group 7</b> containing fungicide before rotating to a fungicide with a different mode of action. Do not apply a third application of this product within 14 days of the second application.</p> <p><b>Restrictions:</b> Do Not apply more than 63 fl oz/A/year. The pre-harvest interval (PHI) for this crop is 40 days. Do Not apply more than the maximum yearly rate from any combination of products containing Isofetamid (1.77 lb. a.i. per acre per year) or Tebuconazole (0.9 lb. a.i. per acre per year) to this crop.</p>

For **24-Hour Medical Emergency Assistance** (Human or Animal) Call 1-888-484-7546.  
For **Chemical Emergency**, Spill, Leak, Fire or Accident, Call CHEMTREC 1-800-424-9300.

Read and follow all applicable directions for use, precautions and limitations on the product label attached to the FERVENT™ 475SC Fungicide container.



Summit Agro USA, LLC  
240 Leigh Farm Road, Suite 215  
Durham, NC 27707