

FERVENT®

475SC FUNGICIDE

Maximize disease control and resistance management by combining two powerful active ingredients

FERVENT® is a valuable tool to help you improve yield and quality in grapes and almonds

FERVENT® 475SC fungicide combines two effective active ingredients to give you an ideal alternative for the control of key almond and grape diseases. FERVENT combines the outstanding preventative disease fighting ability, excellent rainfastness, and translaminar movement of KENJA® fungicide – one of the market's most recent SDHI fungicides – with the proven curative activity of a leading triazole fungicide, tebuconazole.

One reason FERVENT provides this excellent disease control is because of its SDHI fungicide component, which is one of the most active in its class of chemistry. Research indicates that this SDHI has molecular flexibility and because of its unique molecular structure, it has one of the lowest potentials for the development of fungicide resistance.

On grapes, this fungicide combination provides excellent control of Powdery Mildew, along with built-in resistance management through two different fungicide modes of action. The SDHI fungicide component of FERVENT provides additional value through its control of *Botrytis*.

In a disease management program in grapes, the dual modes of action, providing both preventative and curative activity, make FERVENT the right partner to alternate with other Powdery Mildew products, and can fit into any slot in your fungicide program. Or you can start with FERVENT as the first product in your spray program, and alternate it with other Powdery Mildew fungicides, such as PROLIVO® 300SC fungicide. Best of all, FERVENT delivers excellent control of Powdery Mildew on both vines and fruit.

For almond growers, FERVENT is designed to provide excellent curative and preventative control of the key diseases impacting almonds, including Brown rot blossom blight, Anthracnose, Gray Mold, and Shot hole. FERVENT provides effective disease control to meet both early and late season needs.

Registered crops and disease targets:



Grapes

Powdery Mildew
Black Rot
Phomopsis Cane & Leaf Spot
Botrytis Bunch Rot



Almonds

Brown rot blossom blight
Anthracnose
Green fruit rot (*Botrytis*)
Shot hole
Scab
Alternaria leaf spot
Hull rot (suppression)

GROUP 3 FUNGICIDE

GROUP 7 FUNGICIDE

Applying FERVENT

CROP	DISEASES	USE RATE	ADDITIONAL USE INFORMATION
Almond	<p>Brown rot blossom blight (<i>Monilinia spp.</i>)</p> <p>Anthrachnose (<i>Colletotrichum spp.</i>)</p> <p>Green fruit rot (<i>Botrytis cinerea</i>)</p> <p>Shot hole (<i>Wilsonomyces carpophilus</i>)</p> <p>Scab (<i>Cladosporium carpophilum</i>)</p> <p>Alternaria leaf spot (<i>Alternaria alternata</i>)</p> <p>SUPPRESSION:</p> <p>Hull rot (<i>Monilinia spp.</i>, <i>Rhizopus spp.</i>)</p>	12.5 to 15 fl oz per acre	<p>Application Instructions:</p> <p>Initiate applications for brown rot blossom blight when conditions are favorable for disease development and continue on a 7-14 day interval. Typically applications are started preventatively at pink bud and continued through petal fall.</p> <p>Initiate application for control of anthracnose, gray mold, shot hole, scab, Alternaria leaf spot or suppression of hull rot preventatively and continue as needed on a 7-14 day interval. If disease pressure is severe use the higher rate and shortest interval.</p> <p>Apply this product in sufficient water to obtain adequate coverage of the foliage. Spray volume will usually be 50 to 100 gallons per acre for dilute sprays and 5 to 10 gallons per acre for aerial sprays. For aerial applications, apply this product in a minimum of 5 gallons of water per acre. For dilute sprays, if higher spray volumes are desired for improved coverage, do not exceed the maximum rate of 15 fl oz of product per acre.</p> <p>Resistance Management:</p> <p>Do not make more than 2 sequential applications of this product or other Group 3 or Group 7 containing fungicide before rotating to a fungicide with a different mode of action. Do not apply a third application of this product within 14 days of the second application.</p> <p>Restrictions:</p> <p>Do not apply more than 63 fl oz/A/year. The pre-harvest interval (PHI) for this crop is 40 days. Do not apply more than the maximum yearly rate from any combination of products containing Isofetamid (1.77 lb. a.i. per acre per year) or Tebuconazole (0.9 lb. a.i. per acre per year) to this crop.</p>
Crop Subgroup 13-07F (Grapes)	<p>Powdery Mildew (<i>Erysiphe necator</i>)</p> <p>Black rot (<i>Guignardia bidwellii</i>)</p> <p>Phomopsis cane and leaf spot (<i>Phomopsis viticola</i>)</p> <p>Botrytis bunch rot (<i>Botrytis cinerea</i>)</p>	8.5 fl oz per acre	<p>Application Instructions:</p> <p>For use on all types of grapes (wine, table, raisin, and juice).</p> <p>For powdery mildew, black rot, Phomopsis cane and leaf spot, begin fungicide applications preventatively and continue as needed on a 21-day interval. When disease pressure is severe use a shorter interval.</p> <p>For bunch rot make applications at critical timings for Botrytis. Applications are typically made at early bloom, bunch closure, veraison and pre-harvest (at least 14 days apart). Apply with sufficient water to allow for penetration into the foliage to obtain complete coverage using 50 to 100 gallons of spray volume per acre.</p> <p>Resistance Management:</p> <p>Do not make more than 2 sequential applications of this product or other Group 3 or Group 7 containing fungicides before rotating to a fungicide with a different mode of action. Do not apply a third application of this product within 28 days of the second application.</p> <p>Restrictions:</p> <p>Do not apply more than 25.5 fl oz/A/year The Pre-Harvest Interval (PHI) for this crop is 14 days. This product may be applied via aerial application equipment except in the State of New York. Do not apply more than the maximum yearly rate from any combination of products containing Isofetamid (1.72 lb. a.i. per acre per year) or Tebuconazole (0.9 lb. a.i. per acre per year) to these crops.</p>

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of twelve (12) hours. Refer to label for complete use information and restrictions.

FERVENT IS SOLD EXCLUSIVELY THROUGH HELENA AGRI-ENTERPRISES AND TENKOZ MEMBER COMPANIES

LEARN MORE AT [SUMMITAGRO-USA.COM](https://www.summitagro-usa.com)

Always read and follow label directions. FERVENT, PROLIVO and KENJA are inventions and registered trademarks of Ishihara Sangyo Kaisha, Ltd., and are manufactured and developed by ISK Biosciences Corporation. The Summit Agro logo is a trademark of Sumitomo Corporation. All other trademarks are the properties of their respective companies. © 2024 Summit Agro USA LLC.