



TIMOREX ACT[®]

This supplemental label expires June 14, 2024 and must not be used or distributed after this date.

THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THE LABEL AFFIXED TO THE CONTAINER FOR TIMOREX ACT BEFORE APPLYING. USE OF TIMOREX ACT ACCORDING TO THIS LABELING IS SUBJECT TO THE USE PRECAUTIONS AND LIMITATIONS IMPOSED BY THE LABEL AFFIXED TO THE CONTAINER FOR TIMOREX ACT.

ACTIVE INGREDIENT:

Tea Tree Oil 12.5%

OTHER INGREDIENTS: 87.5%

TOTAL: 100.0%

This product contains 0.949 lb. tea tree oil per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you **DO NOT** understand the label, find someone to explain it to you in detail.)

FIRST AID	
If On Skin Or Clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 – 20 minutes. Call a poison control center or doctor for treatment advice.
If Inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Poison Control – National Capital Poison Center 24 hours, 365 days/year	
1-800-222-1222	
NOTE TO PHYSICIAN – No special antidote. Treat symptomatically and supportively.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

See booklet label for additional precautionary statements.

EPA Reg. No.: 86182-3-88783

Manufactured by:

Stockton (Israel) Ltd.
P.O. Box 3517, 17 Ha'Mefalsim St.
Petach Tikva, 4951447, Israel
Tel: +972-72-2570000
www.stk-ag.com
http://stkag.com/product/timorex-act/

Marketed by:

Summit Agro USA, LLC
240 Leigh Farm Rd., Suite 415
Durham, NC 27707
984-260-0407

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION: Harmful if inhaled. Avoid breathing (dust, vapor, or spray mist). Remove and wash contaminated clothing before reuse. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear protective clothing and gloves.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers and loaders must wear:

- Long-sleeved shirt
- Long pants
- Socks with shoes
- Waterproof gloves including barrier laminate, butyl rubber ≥14mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride ≥14 mils, and viton ≥14 mils
- Protective eyewear

Applicators must wear:

- Long-sleeved shirt
- Long pants
- Socks with shoes

Follow manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is harmful to aquatic invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. **DO NOT** contaminate water when cleaning equipment or disposing of equipment wash waters.

PHYSICAL AND CHEMICAL HAZARDS

Combustible. DO NOT use or store near heat or open flame.

DIRECTIONS FOR USE

Use **TIMOREX ACT** on the foliage and fruit of the following agricultural commodities and horticultural crops:

Open Field Crops

Crop	Target Diseases	Fl. Oz. TIMOREX ACT/Acre (lb. ai/acre)	Application Instructions
Pome Fruits: Crop Group 11-10 Apple Azarole Crabapple Loquat Mayhaw Hook & Arn. Medlar Pear Asian Pear Quince Chinese Quince Japanese Quince Tejocote Cultivars, varieties, and/or hybrids of these	Foliar Diseases: Alternaria Blotch (<i>Alternaria mali</i>) Apple Scab (<i>Venturia inaequalis</i>) Bitter Rot (<i>Glomerella cingulata</i> , <i>Colletotrichum</i> spp.) Black Rot, frog-eye leafspot and canker (<i>Botryosphaeria obtusa</i>) Bot Rot (<i>Botryosphaeria dothidea</i>) Brooks Fruit Spot (<i>Mycosphaerella pomi</i>) Bull's Eye Rot (<i>Neofabraea</i> spp.) Flyspeck (<i>Schizothyrum pomi</i>) Moldy Core and Core Rot (<i>Alternaria</i> spp., <i>Cladosporium</i> spp.) Pear Scab (<i>Venturia pirina</i>) Powdery Mildew (<i>Podosphaera leucotricha</i>) White Rot (<i>Botryosphaeria</i> spp.)	14 – 23 (0.104 – 0.169)	Make applications in the early stages of plant growth when conditions favor disease. Use higher rates under increased disease pressure. Re-apply at 7-14 day intervals or as needed throughout the growing season for preventative control. PHI: 14 days.
Stone Fruits: Crop Group 12-12 Apricot Japanese Apricot Capulin Black Cherry Nanking Cherry Sweet Cherry Tart Cherry Chinese Jujube Nectarine Peach Plum American Plum Beach Plum Chanda Plum Cherry Plum Chickasaw Plum Damson Plum Japanese Plum Klamath Plum Prune Plum Plumcot Sloe Cultivars, varieties, and/or hybrids of these	Foliar Diseases: Alternaria Spot/Fruit Rot (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum</i> spp.) Bacterial Canker (<i>Pseudomonas</i> spp.) Bacterial Spot (<i>Pseudomonas</i> spp.) Brown Rot Blossom Blight and Fruit Rot (<i>Monilinia</i> spp.) Cercospora Leaf Spot (<i>Cercospora</i> spp.) Cherry Leaf Spot (<i>Blumeriella jaapii</i>) Gray Mold (<i>Botrytis cinerea</i>) Jacket Rot, Green Fruit Rot (<i>Botrytis cinerea</i> , <i>Monilinia</i> spp., <i>Sclerotinia sclerotiorum</i>) Powdery Mildew (<i>Podosphaera</i> spp., <i>Sphaerotheca pannosa</i>) Rusty Spot (<i>Podosphaera leucotricha</i>) Scab (<i>Cladosporium carpophilum</i>) Shot Hole (<i>Wilsonomyces carpophilus</i>)	14 – 30 (0.104 – 0.221)	Make applications in the early stages of plant growth when conditions favor disease. Use higher rates under increased disease pressure. Re-apply at 7-14 day intervals or as needed throughout the growing season for preventative control up until 48 hours of harvest.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in original container, in a dry, cool place out of direct sunlight and away from heat sources. Keep from overheating or freezing.

Pesticide Disposal: Wastes resulting from this product must be disposed of on site or at an approved waste disposal facility.

Container Handling:

(For containers ≤ 5 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. **DO NOT** burn, unless allowed by State and local ordinances.

(For containers ≥ 30 gallons)

Non-refillable container. **DO NOT** reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in sanitary landfill, or incineration. **DO NOT** burn, unless allowed by State and local ordinances.

WARRANTY STATEMENT

To the extent consistent with the law, seller makes no warranty express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. To the extent consistent with the law, user assumes all risks of use, storage or handling not in strict accordance with label instructions.