



HARVANTA[®] 50SL

INSECTICIDE

EPA Reg. No. 71512-26-88783

Directions for Use on Tuberous and Corm Vegetables (Crop Subgroup 1C) to Control Colorado Potato Beetle in California, Colorado, Florida, Idaho, Maine, Massachusetts, Michigan, Minnesota, Nebraska, New York, North Carolina, North Dakota, Oregon, Pennsylvania, Rhode Island, Virginia, Washington and Wisconsin

PRODUCT BULLETIN

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS

Crop	Insects	Use Rate Fl. Oz. Product Per Acre	Instructions
Tuberous and Corm Vegetables (Crop Subgroup 1C)*	Colorado potato beetle	5.5 to 10.9 fl oz (0.018 to 0.036 lb. a.i. /A)	<p>Application Instructions: Thorough coverage is essential to achieve best results.</p> <p>For Colorado Potato Beetle: Use the higher application rate under heavy pest pressure and/or when larger larvae or adults are present</p> <p>Resistance Management: Do not apply HARVANTA 50SL INSECTICIDE or other Group 28 insecticide more than 3 times within a single generation of insect pest(s) on a crop.</p> <p>Restrictions: Foliar application of this product is prohibited to a crop from onset of flowering until flowering is complete unless:</p> <ul style="list-style-type: none"> • The application is being made in the time period between 2 hours prior to sunset until sunrise; OR, • The application is being made at a time when the temperature at the application site is 50 degrees F or less. <p>[In California no more than one application may be made from the onset of flowering until flowering is complete. Applications must be made when bees are not foraging, adhering to the above restrictions.]</p> <p>Make no more than 3 applications per year. Minimal interval between treatment is 5 days [14 days in California]. Do not apply more than 49.2 fl oz per acre per year (0.16 lb. a.i./acre/year). Pre-Harvest Interval (PHI) – 7 days</p>

*Includes all members of the Tuberous and Corm Vegetable Crop Group Family: Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; taniel; turmeric; yam bean; yam, true

For **24-Hour Medical Emergency Assistance** (Human or Animal) Call 1-888-484-7546.
For **Chemical Emergency**, Spill, Leak, Fire or Accident, Call CHEMTREC 1-800-424-9300.

Read and follow all applicable directions for use, precautions and limitations on the product label attached to the Harvanta 50SL Insecticide container.



Summit Agro USA, LLC
240 Leigh Farm Road, Suite 415
Durham, NC 27707

Harvanta is an invention and registered trademark of Ishihara Sangyo Kaisha, Ltd., and is manufactured and developed by ISK Biosciences Corporation.



HARVANTA[®] 50SL

INSECTICIDE

Expires 5/20/2024

EPA Reg. No. 71512-26-88783

Directions for Use on Tuberous and Corm Vegetables (Crop Subgroup 1C) to Control Colorado Potato Beetle in the State of Texas only

PRODUCT BULLETIN

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the EPA. This product bulletin should be in the possession of the user at the time of application of the product. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

APPLICATION DIRECTIONS

Crop	Insects	Use Rate Fl. Oz. Product Per Acre	Instructions
Tuberous and Corm Vegetables (Crop Subgroup 1C)*	Colorado potato beetle	5.5 to 10.9 fl oz (0.018 to 0.036 lb. a.i. /A)	<p>Application Instructions: Thorough coverage is essential to achieve best results.</p> <p>For Colorado Potato Beetle: Use the higher application rate under heavy pest pressure and/or when larger larvae or adults are present</p> <p>Resistance Management: Do not apply HARVANTA 50SL INSECTICIDE or other Group 28 insecticide more than 3 times within a single generation of insect pest(s) on a crop.</p> <p>Restrictions: Foliar application of this product is prohibited to a crop from onset of flowering until flowering is complete unless:</p> <ul style="list-style-type: none"> • The application is being made in the time period between 2 hours prior to sunset until sunrise; OR, • The application is being made at a time when the temperature at the application site is 50 degrees F or less. <p>[In California no more than one application may be made from the onset of flowering until flowering is complete. Applications must be made when bees are not foraging, adhering to the above restrictions.]</p> <p>Make no more than 3 applications per year. Minimal interval between treatment is 5 days [14 days in California]. Do not apply more than 49.2 fl oz per acre per year (0.16 lb. a.i./acre/year). Pre-Harvest Interval (PHI) – 7 days</p>

*Includes all members of the Tuberous and Corm Vegetable Crop Group Family: Arracacha; arrowroot; artichoke, Chinese; artichoke, Jerusalem; canna, edible; cassava, bitter and sweet; chayote (root); chufa; dasheen (taro); ginger; leren; potato; sweet potato; taniel; turmeric; yam bean; yam, true

For **24-Hour Medical Emergency Assistance** (Human or Animal) Call 1-888-484-7546.
For **Chemical Emergency**, Spill, Leak, Fire or Accident, Call CHEMTREC 1-800-424-9300.

Read and follow all applicable directions for use, precautions and limitations on the product label attached to the Harvanta 50SL Insecticide container.



Summit Agro USA, LLC
240 Leigh Farm Road, Suite 415
Durham, NC 27707

Harvanta is an invention and registered trademark of Ishihara Sangyo Kaisha, Ltd., and is manufactured and developed by ISK Biosciences Corporation.